

# Proceedings of the Iowa Academy of Science

---

Volume 39 | Annual Issue

Article 56

---

1932

## Resistivity of Zinc Crystals

W.J. Poppy

*State University of Iowa*

Copyright © Copyright 1932 by the Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Poppy, W. J. (1932) "Resistivity of Zinc Crystals," *Proceedings of the Iowa Academy of Science*, 39(1), 217-217.

Available at: <https://scholarworks.uni.edu/pias/vol39/iss1/56>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

## A NEW VACCUUM TYPE MERCURY STILL

L. E. PINNEY

The still consists of a compact single unit of Pyrex glass embodying a boiler, a water cooled condenser, and a receiver. Mercury can be poured from the receiver as required without disturbing the contents of the boiler.

DEPARTMENT OF PHYSICS,  
IOWA STATE COLLEGE,  
AMES, IOWA.

---

## THERMOELECTRIC POWER IN BISMUTH SINGLE CRYSTALS IN THE NEIGHBORHOOD OF THE MELTING POINT

A. SOROOS

The thermoelectric power is determined directly and studied as a function of temperature for a range extending from several degrees below the melting point to a temperature well above the melting point. A transition region exists for about ten degrees above the melting point. This is taken as a definite indication of the persistence of a marked crystalline arrangement in the liquid near its freezing point.

DEPARTMENT OF PHYSICS,  
STATE UNIVERSITY OF IOWA,  
IOWA CITY, IOWA.

---

## RESISTIVITY OF ZINC CRYSTALS

W. J. POPPY

In an attempt to settle the discrepancy between the resistivity measurements of Bridgman on the one hand and Tyndall and Hoyem (Phys. Rev. 38, p. 820; 1931) on the other, single zinc crystals of one square cm. cross section and ten cm. long have been grown and measured. The results are in agreement with Tyndall and Hoyem. Present indications are that certain **anomalous** crystals (i.e. not truly single) have abnormally low resistivities and show great sensitivity to slight strain. The technique of growing these large crystals will be described.

DEPARTMENT OF PHYSICS,  
STATE UNIVERSITY OF IOWA,  
IOWA CITY, IOWA.